

1    **Chemotherapy induced intestinal mucosal barrier damage: a cause of falsely elevated**  
2    **serum 1,3-beta-d-glucan levels?**

3

4    Juergen Prattes<sup>1,2</sup>, Reinhard B. Raggam<sup>3,\*</sup>, Kim Vanstraelen<sup>4</sup>, Jasmin Rabensteiner<sup>3</sup>,  
5    Christoph Hoegenauer<sup>5</sup>, Robert Krause<sup>1,2</sup>, Florian Prueller<sup>3</sup>, Albert Woelfler<sup>6</sup>, Isabel Spriet<sup>4</sup>,  
6    Martin Hoenigl<sup>1,2,7,8,#</sup>

7    1 Section of Infectious Diseases and Tropical Medicine, Medical University of Graz, Graz,  
8    Austria

9    2 Center for Biomarker Research in Medicine, Graz, Austria

10    3 Clinical Institute of Medical and Chemical Laboratory Diagnostics, Medical University of  
11    Graz, Graz, Austria

12    4 Clinical Pharmacology and Pharmacotherapy, Department of Pharmaceutical and  
13    Pharmacological Sciences University Hospital Leuven, Leuven, Belgium

14    5 Division of Gastroenterology and Hepatology, Medical University of Graz, Graz, Austria

15    6 Division of Hematology, Medical University of Graz, Graz, Austria

16    7 Division of Infectious Diseases, Department of Medicine, University of California–San  
17    Diego, San Diego, USA

18    8 Division of Pulmonology, Medical University of Graz, Graz, Austria

19

20

21    Running Head: Serum BDG levels in intestinal mucositis

22

23 #Address correspondence to Martin Hoenigl, M.D., [martin.hoenigl@medunigraz.at](mailto:martin.hoenigl@medunigraz.at) or

24 [mhoenigl@ucsd.edu](mailto:mhoenigl@ucsd.edu)

25 Present address: Division of Infectious Diseases, Department of Medicine, University of

26 California, San Diego,

27 200 West Arbor Drive #8208 San Diego, CA 92103, United States of America

28 Phone: +1 6195435605

29

30 \*Alternative corresponding author: Reinhard B. Raggam, M.D.,

31 [reinhard.raggam@medunigraz.at](mailto:reinhard.raggam@medunigraz.at)

32 Present address: Clinical Institute of Medical and Chemical Laboratory Diagnostics, Medical

33 University of Graz, Graz, Austria, Auenbruggerplatz 15, 8036 Graz, Austria

34 Phone: +43 31638580243

35

36 Abstract

37

38 Blood citrulline and intestinal fatty acid binding protein were determined as biomarkers for  
39 intestinal mucositis. Biomarker levels were correlated with corresponding serum 1,3-beta-d-  
40 glucan levels in 56 samples obtained from 33 cases with underlying hematological  
41 malignancies receiving induction chemotherapy. No correlation between biomarkers of  
42 intestinal mucositis and BDG levels was observed.

43

44 (1→3)-β-D-Glucan (BDG) is a polysaccharide cell wall component of most fungal species  
45 and a useful serum biomarker for early diagnosis of invasive fungal infections (IFIs) (1, 2).  
46 False positive results have been linked to use of cellulose containing haemodialysis  
47 membranes, blood-derivates, broad spectrum antibiotics, and severe mucositis (3). Intestinal  
48 mucositis is a frequently observed adverse event of induction chemotherapy in patients with  
49 underlying hematological malignancies (4). Severe mucositis is characterized by a loss of  
50 integrity of the intestinal mucosal barrier, increasing the likelihood of translocation of  
51 bacterial and/or fungal commensals of the gastrointestinal tract (5). Whether intestinal  
52 mucositis is a cause of false positive BDG serum levels has been insufficiently evaluated to  
53 date mainly because of difficulties in diagnosing and defining mucositis in hematological  
54 malignancy patients in clinical routine. Biopsy of the gastrointestinal tract is the diagnostic  
55 gold-standard, but often hindered by the unfavorable risk-benefit ratio in those with severe  
56 underlying conditions and increased risk of bleeding.

57 Thus, several clinical scores have been implemented for diagnosis and grading of mucositis  
58 (6, 7), but these are prone to subjective interpretation. Therefore, noninvasive diagnostic  
59 biomarkers for mucositis have been increasingly investigated. Serum citrulline, a non-protein  
60 amino acid produced by intact enterocytes (8, 9), and intestinal fatty acid binding protein  
61 (IFABP), low-molecular mass cytosolic proteins found in tissues involved in the uptake,  
62 intracellular metabolism and transport of long-chain fatty acids (10, 11), are two of the most  
63 promising biomarkers for detection of loss of enterocytes and therefore mucosal barrier  
64 injury.

65 The objective of this study was to investigate whether serum BDG levels are falsely elevated  
66 in hematological malignancy patients with intestinal mucositis. A total of 56 same day plasma  
67 and serum samples, were prospectively obtained from 33 consecutive cases (i.e. 33  
68 admissions corresponding to 24 patients; maximum 2 samples per case) with underlying

69 hematological malignancies receiving induction chemotherapy for acute leukemia. None of  
70 our patients fulfilled criteria for IFI according to 2008 revised definitions for IFIs from the  
71 EORTC/MSG group (12, 13). Samples were collected between July 2012 and May 2013 at  
72 the University Hospital of Graz, Austria, and a maximum of two samples per case were  
73 included (in those with two samples, samples were obtained always > 4 days apart). BDG was  
74 determined prospectively, while citrulline and IFABP were determined retrospectively in  
75 serum/plasma samples after immediate sample storage at -70° at the Medical University of  
76 Graz, Austria. BDG was determined using the automated protocol Fungitell® assay  
77 (Associates of Cape Cod, Inc., East Falmouth, USA), as described previously (14). IFABP  
78 was determined using the Quantikine Human FABP2/I-FABP Immunoassay (R&D Systems  
79 Europe Ltd., OX, UK) according to the manufacturer's instruction. Citrulline was determined  
80 from plasma samples at the University Hospital Leuven, Belgium, using liquid  
81 chromatograph-tandem mass spectrometry as described elsewhere (15). Mucositis was  
82 classified according to citrulline levels: >30µmol/L as no evidence of mucositis, 20-30µmol/L  
83 as partial villus atrophy, 10-20µmol/L as destructive mucosal lesions and <10µmol/L as  
84 diffuse destructive mucositis (16, 17). IFABP was used as a second marker for mucositis to  
85 better support mucositis classification based on citrulline levels. In contrast to citrulline  
86 IFABP levels are highly elevated in case of intestinal mucosal barrier damage, with 100 fold  
87 increases in patients with intestinal ischemia when compared to healthy controls (10).

88 For statistical analyses a cut-off of 10µmol/L citrulline was used as indicator for extensive  
89 mucositis. The study adhered to the Declaration of Helsinki (1996) and Good Clinical  
90 Practice. The study protocol was approved by the local ethics committee, Medical University  
91 Graz, Austria (EC-number 23-343) and registered at ClinicalTrials.gov (identifier  
92 NCT01576653). Informed consent was obtained from all participating patients. For statistical  
93 analysis SPSS Version 23 was used (SPSS Inc., Chicago, IL, USA). For continuous variables

94 medians plus interquartile range (IQR) or mean plus 95% confidence interval (95% CI) are  
95 displayed as appropriate. Correlation analysis was performed by using Spearman-Rho  
96 analyses due to non-normality of our data and data are presented in scatterplots. Retrospective  
97 power analysis revealed that a sample size of 48 mucositis samples (8 samples were derived  
98 from patients without mucositis) provides at least 80% power (with  $\alpha = 0.05$ ) to detect a  
99 correlation of  $r=0.44$  (i.e.  $r^2=0.19$ ) or higher for correlations of three variables. Chi-squared  
100 test was used for categorical data. A p-value of  $<0.05$  was considered statistically significant.

101 Demographic data, underlying diseases, and biomarker levels for patients, cases and samples  
102 are displayed in Table1. All patients were receiving mold-active antifungal prophylaxis with  
103 posaconazole oral suspension, as recommended in clinical guidelines. Additionally, all  
104 included patients were able to tolerate solid food. All but two patients were on systemic  
105 antibiotic treatment/prophylaxis at the time of sample collection. Most common antibiotics  
106 prescribed at the time of sampling were: levofloxacin (23 samples, 42.6%), cefepime (21  
107 samples, 38.9%), linezolid (7 samples, 13%) and meropenem (6 samples, 11.1%).  
108 Additionally, all patient received prophylaxis for oral mucositis containing chlorhexidine  
109 mouth wash and nystatin oral solution. Overall specificity of serum BDG testing in our cohort  
110 was 0.95 (95% CI 0.85 – 0.98).

111 Positive BDG serum levels ( $>80\text{pg/ml}$ ) were observed in three serum samples: two out of 17  
112 samples (11.8%) with citrulline levels  $<10\mu\text{mol/L}$  had false positive BDG levels (120.5pg/ml  
113 and 149.3pg/ml respectively; corresponding IFABP levels: 1366.3pg/ml and 976.1pg/ml  
114 respectively) and one out of 39 samples (2.6%) with citrulline levels  $\geq 10\mu\text{mol/L}$  had a false  
115 positive BDG level (232.4pg/ml; corresponding IFABP level: 380.1pg/ml;  $p=0.160$ ). Median  
116 BDG levels in samples with citrulline levels  $<10\mu\text{mol/L}$  and  $\geq 10\mu\text{mol/L}$  were 15.4pg/ml in  
117 both groups. Serum IFABP levels showed a strong negative correlation with plasma citrulline  
118 levels (Spearman-Rho  $r=-0.601$ ;  $p=<0.001$ ). Additionally, IFABP values were significantly

119 higher in the mucositis group (i.e. n=17 with  $<10\mu\text{mol/L}$  citrulline) compared to the group  
120 without severe mucositis according to citrulline levels (814.8pg/ml versus 380.9pg/ml;  
121  $p=0.004$ ) supporting mucositis diagnosis based on citrulline levels. Correlations between  
122 BDG and citrulline, or IFABP are displayed in Figure 1.

123 Our main objective in this study was to determine whether underlying intestinal mucositis in  
124 hematological malignancy patients is a potential cause of falsely elevated serum BDG levels.  
125 As surrogate markers for mucositis we used citrulline and IFABP. Both markers showed  
126 excellent correlation with each other supporting the diagnosis of intestinal mucositis,  
127 however, no correlation between either citrulline or IFABP and serum BDG levels was  
128 observed. Additionally, only 2 out of 17 samples from patients with biomarker levels  
129 suggestive for diffuse intestinal mucosal lesion yielded positive serum BDG levels.

130 As fungi, especially *Candida spp.*, are commensals in the gastrointestinal tract, mucosal  
131 barrier damage is considered a risk factor for developing IFIs (18, 19). Antifungal prophylaxis  
132 is therefore a recommended practice in patients with hematological malignancies receiving  
133 induction chemotherapy or hematopoietic stem cell transplantation (20). In our setting oral  
134 posaconazole suspension was used as prophylactic agent in all patients. Antifungal  
135 prophylaxis is likely to result in a decreased fungal burden in the gastrointestinal tract and  
136 therefore a reduced amount of fungal components in serum in case of mucosal barrier  
137 damage. This may be particularly true for the posaconazole suspension that was used in all  
138 our patients. Due to the liquid formulation, posaconazole suspension is acting more fungicidal  
139 in the upper gastrointestinal tract (i.e. oral, esophagus) when compared to the newer tablet  
140 formulation. As fungal colonization is mainly found in the upper gastrointestinal tract,  
141 posaconazole suspension may cause more substantial reduction of fungal colonization when  
142 compared to the tablet or the intravenous form. While the fact that antifungal prophylaxis was  
143 used in all participants may be considered a limitation from a scientific standpoint (i.e. we

144 could not evaluate whether mucositis is associated with false positive BDG levels in the  
145 absence of antifungal prophylaxis), it may be considered as strength from a clinical standpoint  
146 as this reflects clinical practice in most specialized centers.

147 Specificity of serum BDG testing in our study was 95% which is comparable to other studies  
148 in hematological malignancy patients (21). Our results therefore clearly indicate that  
149 mucositis may not be a major cause of false positive BDG levels in patients with induction  
150 chemotherapy, receiving antifungal prophylaxis. These findings have important clinical  
151 implications, as false negative BDG test results in the presence of antifungal therapy or  
152 prophylaxis have been shown to occur rarely. This is supported by a study conducted by Koo  
153 et al., who could demonstrate that systemic antifungal therapy did not alter the performance of  
154 serum BDG determination (22).

155 Our study was limited by indirect assessment of intestinal mucositis by citrulline and IFABP  
156 determination. No endoscopies to verify intestinal mucosal barrier damage were performed.  
157 Additionally, citrulline and IFABP are both produced in the lower gastrointestinal tract, and  
158 may not be very reliable biomarkers in patients with exclusively oral mucositis. Oral  
159 mucositis may have therefore been underestimated in our study. Small samples size was  
160 another important limitation.

161 In conclusion, we did not observe elevated serum BDG levels in hematological malignancy  
162 patients with underlying intestinal mucositis who received antifungal prophylaxis, indicating  
163 that serum BDG may be a specific biomarker in this setting.

164

165

166

167

168



169 **Acknowledgements**

170 Data of the manuscript have been previously presented, in part, at the ICAAC, San Diego  
171 2015 (guided poster walk, presentation number 1477) and at the TIMM, Lisbon 2015  
172 (presentation number P452).

173

174 **Conflicts of interest**

175 Reinhard B. Raggam received travel grants from Pfizer and Merck.

176 A.Wölfler received research grants and speaker honoraria from Merck.

177 M. Hoenigl received research grants from Merck and Pfizer; served on the speakers' bureau  
178 of Pfizer, Gilead, Astellas, Basilea and Merck and received travel grants from Astellas,  
179 Merck, Gilead and Pfizer.

180 I. Spriet received research grants from Merck and Pfizer and speaker honoraria and travel  
181 grants from Pfizer, Merck and Gilead.

182 All other authors no conflict.

183

184 **Funding**

185 This work was supported by funds of the Oesterreichische Nationalbank (Anniversary Fund,  
186 project number 15346), and a Merck Investigator Studies Program (39543) research grant.

187 The funders had no role in study design, data collection, analysis, interpretation, decision to  
188 publish, in the writing of the manuscript, and in the decision to submit the manuscript for  
189 publication.

## References

1. Karageorgopoulos DE, Vouloumanou EK, Ntziora F, Michalopoulos A, Rafailidis PI, Falagas ME. 2011. beta-D-glucan assay for the diagnosis of invasive fungal infections: a meta-analysis. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **52**:750-770.
2. Prattes J, Hoenigl M, Rabensteiner J, Raggam RB, Pruessner F, Zollner-Schwetz I, Valentin T, Hoenigl K, Fruhwald S, Krause R. 2014. Serum 1,3-beta-d-glucan for antifungal treatment stratification at the intensive care unit and the influence of surgery. *Mycoses*.
3. Ellis M, Al-Ramadi B, Finkelman M, Hedstrom U, Kristensen J, Ali-Zadeh H, Klingspor L. 2008. Assessment of the clinical utility of serial beta-D-glucan concentrations in patients with persistent neutropenic fever. *Journal of medical microbiology* **57**:287-295.
4. Benson AB, 3rd, Ajani JA, Catalano RB, Engelking C, Kornblau SM, Martenson JA, Jr., McCallum R, Mitchell EP, O'Dorisio TM, Vokes EE, Wadler S. 2004. Recommended guidelines for the treatment of cancer treatment-induced diarrhea. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* **22**:2918-2926.
5. Bounoux ME, Diogo D, Francois N, Sendid B, Veirmeire S, Colombel JF, Bouchier C, Van Kruiningen H, d'Enfert C, Poulain D. 2006. Multilocus sequence typing reveals intrafamilial transmission and microevolutions of *Candida albicans* isolates from the human digestive tract. *Journal of clinical microbiology* **44**:1810-1820.
6. Donnelly JP, Muus P, Schattenberg A, De Witte T, Horrevorts A, DePauw BE. 1992. A scheme for daily monitoring of oral mucositis in allogeneic BMT recipients. *Bone marrow transplantation* **9**:409-413.
7. Potting CM, Blijlevens NA, Donnelly JP, Feuth T, Van Achterberg T. 2006. A scoring system for the assessment of oral mucositis in daily nursing practice. *European journal of cancer care* **15**:228-234.
8. Lutgens LC, Deutz N, Granzier-Peters M, Beets-Tan R, De Ruyscher D, Gueulette J, Cleutjens J, Berger M, Wouters B, von Meyenfeldt M, Lambin P. 2004. Plasma citrulline concentration: a surrogate end point for radiation-induced mucosal atrophy of the small bowel. A feasibility study in 23 patients. *International journal of radiation oncology, biology, physics* **60**:275-285.
9. Barzal JA, Szczylik C, Rzepecki P, Jaworska M, Anuszevska E. 2014. Plasma citrulline level as a biomarker for cancer therapy-induced small bowel mucosal damage. *Acta biochimica Polonica* **61**:615-631.
10. Lieberman JM, Sacchettini J, Marks C, Marks WH. 1997. Human intestinal fatty acid binding protein: report of an assay with studies in normal volunteers and intestinal ischemia. *Surgery* **121**:335-342.
11. Paulussen RJ, Veerkamp JH. 1990. Intracellular fatty-acid-binding proteins. Characteristics and function. *Sub-cellular biochemistry* **16**:175-226.
12. De Pauw B, Walsh TJ, Donnelly JP, Stevens DA, Edwards JE, Calandra T, Pappas PG, Maertens J, Lortholary O, Kauffman CA, Denning DW, Patterson TF, Maschmeyer G, Bille J, Dismukes WE, Herbrecht R, Hope WW, Kibbler CC, Kullberg BJ, Marr KA, Munoz P, Odds FC, Perfect JR, Restrepo A, Ruhnke M, Segal BH, Sobel JD, Sorrell TC, Viscoli C, Wingard JR, Zaoutis T, Bennett JE, European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group, National Institute of Allergy and Infectious Diseases Mycoses Study Group Consensus G. 2008. Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **46**:1813-1821.

- 241 13. **Hoenigl M, Strenger V, Buzina W, Valentin T, Koidl C, Wolfler A, Seeber K, Valentin A,**  
242 **Strohmeier AT, Zollner-Schwetz I, Raggam RB, Urban C, Lass-Flörl C, Linkesch W, Krause R.**  
243 2012. European Organization for the Research and Treatment of Cancer/Mycoses Study  
244 Group (EORTC/MSG) host factors and invasive fungal infections in patients with  
245 haematological malignancies. *The Journal of antimicrobial chemotherapy* **67**:2029-2033.
- 246 14. **Pruller F, Wagner J, Raggam RB, Hoenigl M, Kessler HH, Truschnig-Wilders M, Krause R.**  
247 2014. Automation of serum (1 $\rightarrow$ 3)-beta-D-glucan testing allows reliable and rapid  
248 discrimination of patients with and without candidemia. *Medical mycology* **52**:455-461.
- 249 15. **Demacker PN, Beijers AM, van Daal H, Donnelly JP, Blijlevens NM, van den Ouweland JM.**  
250 2009. Plasma citrulline measurement using UPLC tandem mass-spectrometry to determine  
251 small intestinal enterocyte pathology. *Journal of chromatography. B, Analytical technologies*  
252 *in the biomedical and life sciences* **877**:387-392.
- 253 16. **Crenn P, Messing B, Cynober L.** 2008. Citrulline as a biomarker of intestinal failure due to  
254 enterocyte mass reduction. *Clinical nutrition* **27**:328-339.
- 255 17. **Crenn P, Vahedi K, Lavergne-Slove A, Cynober L, Matuchansky C, Messing B.** 2003. Plasma  
256 citrulline: A marker of enterocyte mass in villous atrophy-associated small bowel disease.  
257 *Gastroenterology* **124**:1210-1219.
- 258 18. **Pittet D, Monod M, Suter PM, Frenk E, Auckenthaler R.** 1994. Candida colonization and  
259 subsequent infections in critically ill surgical patients. *Annals of surgery* **220**:751-758.
- 260 19. **Nucci M, Anaissie E.** 2001. Revisiting the source of candidemia: skin or gut? *Clinical infectious*  
261 *diseases : an official publication of the Infectious Diseases Society of America* **33**:1959-1967.
- 262 20. **Cornely OA, Maertens J, Winston DJ, Perfect J, Ullmann AJ, Walsh TJ, Helfgott D,**  
263 **Holowiecki J, Stockelberg D, Goh YT, Petrini M, Hardalo C, Suresh R, Angulo-Gonzalez D.**  
264 2007. Posaconazole vs. fluconazole or itraconazole prophylaxis in patients with neutropenia.  
265 *The New England journal of medicine* **356**:348-359.
- 266 21. **Marchetti O, Lamothe F, Mikulska M, Viscoli C, Verweij P, Bretagne S, European Conference**  
267 **on Infections in Leukemia Laboratory Working G.** 2012. ECIL recommendations for the use  
268 of biological markers for the diagnosis of invasive fungal diseases in leukemic patients and  
269 hematopoietic SCT recipients. *Bone marrow transplantation* **47**:846-854.
- 270 22. **Koo S, Bryar JM, Page JH, Baden LR, Marty FM.** 2009. Diagnostic performance of the (1 $\rightarrow$ 3)-  
271 beta-D-glucan assay for invasive fungal disease. *Clinical infectious diseases : an official*  
272 *publication of the Infectious Diseases Society of America* **49**:1650-1659.

273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293

294

295

296

297

298

299

300

301

302 **Figure 1 Correlation of 1,3-beta-d-glucan (BDG) with citrulline and intestinal fatty acid**

303 **binding protein (IFABP) in same day plasma and serum samples.** Correlation analyses

304 were performed by Spearman-Rho-analysis and yielded no significant correlations between

305 BDG and citrulline ( $r=0.081$ ) and BDG and IFABP ( $r=0.048$ ).

306

**Table 1** Demographic data, underlying diseases and 1,3-beta-d-glucan (BDG), citrulline and intestinal fatty acid binding protein (IFABP) levels in the study population.

	Patients* [n (%)]	Cases** [n (%)]	Samples*** [n (%)]
N	24	33	56
Sex: Female / Male	10 (41.7) / 14 (58.3)	15 (45.5) / 18 (54.5)	28 (50) / 28 (50)
Age, years (range)	54 (20-67)	57 (20-67)	59 (20-67)
Underlying diseases			
Acute myeloid leukemia	18 (75)	26 (78.8)	46 (82.1)
Acute lymphoblastic leukemia	3 (12.5)	4 (12.1)	5 (8.9)
Myelodysplastic syndrome	2 (8.3)	2 (6.1)	3 (5.4)
Aplastic anemia	1 (4.2)	1 (3)	2 (3.6)
Citrulline $\mu\text{mol/L}$ , median (IQR)	14.3 (8.6-33.6)	14.1 (8.8-18.2)	13.8 (8.7-18.2)
Citrulline $<10\mu\text{mol/L}$ (diffuse destructive mucositis)	8 (33.3)	11 (33.3)	17 (30.4)
Citrulline $10\text{--}20\mu\text{mol/L}$ (destructive mucosal lesions)	12 (50)	18 (54.5)	31 (55.4)
Citrulline $20\text{--}30\mu\text{mol/L}$ (partial villus atrophy)	1 (4.2)	1 (3)	4 (7.1)
Citrulline $\geq 30\mu\text{mol/L}$ (no evidence of mucositis)	3 (12.5)	3 (9.1)	4 (7.1)
BDG pg/ml, median (IQR)	15.4 (7.4-39.9)	15.4 (5-35.8)	15.4 (6.8-30.9)
BDG $\geq 80\text{pg/ml}$	2 (8.3)	2 (6.1)	3 (5.4)
BDG $< 80\text{pg/ml}$	22 (91.7)	31 (93.9)	53 (94.6)
IFABP pg/ml, median (IQR)	524.6 (309.8-804.3)	483.5 (332.3-849.7)	514.9 (337.9-792.6)
Severe Diarrhea	3 (12.5)	3 (9.1)	4 (7.1)
Citrulline $<10\mu\text{mol/L}$ (diffuse)	1 (33.3)	1 (33.3)	2 (50)

destructive mucositis) Citrulline 10-20μmol/L (destructive mucosal lesions)	2 (66.6)	2 (66.6)	2 (50)
Absolute numbers plus percentage displayed, unless otherwise noted. *Patients = first sample per patient within the study period; **Cases = 33 admissions corresponding to 24 patients; ***Samples = all samples obtained from included patients within the study period Abbreviations: BDG = 1,3-beta-d-glucan; IQR = 25 -75 interquartile range; IFABP = intestinal fatty acid binding protein			

